



Tri-Service Post-Operative Rehabilitation Guidelines

May 2020

Approved by:

B. KYLE POTTER

COL, MC, USA

Army Orthopedics Consultant

GEORGE P. NANOS III

CAPT, MC, USN

Navy Orthopedics Consultant

JOSEPH J. STUART

Lt Col, USAF, MC

Air Force Orthopedics Consultant

JASON L. SILVERNAIL

COL, SP, USA

Army Physical Therapy Consultant

LESLIE C. HAIR

CDR, MSC, USN

Navy Physical Therapy Consultant

JAMES E. SHIELDS

Col, USAF, BSC

Air Force Physical Therapy Consultant

Created with support from:



Uniformed Services University of the
Health Sciences –
F. Edward Hébert School of Medicine



The Geneva Foundation



Military Orthopaedics Tracking
Injuries and Outcomes Network
(MOTION)

This work was supported by the Uniformed Services University, Department of Physical Medicine & Rehabilitation, Musculoskeletal Injury Rehabilitation Research for Operational Readiness (MIRROR) (HU00011920011).

The opinions and assertions expressed herein are those of the author(s) and do not necessarily reflect the official policy or position of the Uniformed Services University or the Department of Defense.



Arthroscopic Shoulder Rehabilitation

(Subacromial Decompression, Distal Clavicle Resection, or Debridement)

These guidelines were created as a framework for the post-operative rehabilitation program. They DO NOT substitute for any specific restrictions or requirements that are determined through the necessary shared decision-making and collaboration between the operating surgeon and treating rehabilitation team.

PHASE 1: Generally 0-4 Weeks Post-Op

GOALS:	1) Normal shoulder ROM 2) Pain free ADLs 3) Minimize pain and swelling
PRECAUTIONS:	- <u>NO</u> push-ups, heavy lifting, or other sports participation - <u>NO</u> repetitive overhead use of shoulder - If open DCE, avoid cross body ADD and axial traction x 4 weeks - Pain should be no more than mild to moderate during exercises. Pain should settle relatively quickly and should not inhibit exercises the following day.
SLING:	- Sling use for comfort. Recommended use: • Days 1-3: wear sling ~75% of the time • Days 4-7: wear sling ~50% of the time • Days 8-10: D/C sling
WOUND:	- Post-op dressing removed at PT eval - May shower at post-op day #3 - Submerge in water <u>after</u> wound is fully healed - Suture removal @ 7-14 days post-op by Ortho
MODALITIES:	- Cryotherapy • Hourly for 15 minutes for the first 24 hours <u>after</u> sensation is restored from nerve block • Continue use until acute inflammation is controlled • Once controlled, use 3x per day for 15 minutes or longer as tolerated - Soft tissue mobilization and other integrative medicine techniques • Soft tissue/trigger point work to the kinetic chain (i.e. cervical spine, scapula, and forearm)
REHABILITATION:	<u>NOTE:</u> As tolerated, progress rehabilitation exercises as wound healing occurs and the inflammatory response decreases - ROM exercises - Scapular strengthening emphasizing scapular retractors/upward rotators

	- Shoulder strength and endurance progression as ROM is normalized • Continue base strengthening/isometrics as needed • Consider Blood Flow Restriction Therapy to non-operative and/or operative side as tolerated • Rotator cuff progressive resistance exercises (PREs) • Increase functional activities - Modalities as required - Aerobic conditioning - Adjunct treatments to consider: dry needling, cervicothoracic manual therapy, aquatic therapy, etc.
FOLLOW-UP:	- Supervised rehab: 1-2x per week - PT re-eval: every 2 weeks - Ortho re-eval: ~2 weeks and upon achievement of PT goals
MISCELLANEOUS:	- Meet occupational requirements at 3-6 months - Pass Service fitness test at 6 months - Progress activities for return to sport/collision sports or aggressive military training (i.e. airborne school) based on the patient's functional performance and endurance. This time period will be directed by the Ortho Surgeon and the Physical Therapist. This may require between 4-9 months before cleared without restrictions.

References:

- Orthopaedic Specialists of North Carolina Protocol: Gall and Kirby
- Duke University Arthroscopic Subscapular Decompression Post-Surgical Rehabilitation Protocol
- Massachusetts General Hospital Postoperative Shoulder Rehabilitation Protocols